

Letter to the Editor

Open Access

Antiviral therapy in acute viral hepatitis B: why and when

Giuseppe Morelli¹, Alessandro Perrella*^{1,2}, Costanza Sbreglia¹,
Pasquale Bellopede¹, Vincenzo Riccio¹ and Oreste Perrella¹

Address: ¹VII Department of Infectious Diseases and Immunology, "D. Cotugno" Hospital, Naples, Italy and ²"Liver Unit," A. Cardarelli Hospital, Naples, Italy

Email: Giuseppe Morelli - pinomorelli25@libero.it; Alessandro Perrella* - alexperrel@virgilio.it; Costanza Sbreglia - drcosty@libero.it; Pasquale Bellopede - pbellop@libero.it; Vincenzo Riccio - oreste.perrella@ospedalecotugno.it; Oreste Perrella - oreste.perrella@ospedalecotugno.it

* Corresponding author

Published: 16 January 2009

Received: 30 July 2008

Accepted: 16 January 2009

Infectious Agents and Cancer 2009, **4**:2 doi:10.1186/1750-9378-4-2

This article is available from: <http://www.infectagentscancer.com/content/4/1/2>

© 2009 Morelli et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Acute viral hepatitis B is cleared in more than 95% of patients, while the remainder ones may develop either chronic HBV infection or, rarely, fulminant hepatitis.

Therefore there are elderly patients with severe acute HBV hepatitis characterized by high serum bilirubin levels >15 mmole/dl, international normalized ratio (INR) with value more than 1.6; these patients are characterized by a severe outcome of HBV infection.

As known, outcome of infection and the pathogenesis of liver diseases are determined by viral and host factors, such as T reg lymphocytes.

T regs may be associated with a negative immune response such as an inhibition of gamma- IFN secretion.

The impact of viral load on antiviral T cell responses may play a critical role in these patients, influencing disease persistence and immune response.

Antiviral drugs could be useful in these patients determining a possible down -regulation of T regs.

Letter to the editor

Dear Editor,

Acute viral hepatitis B is cleared in more than 95% of patients, while the remainder ones may develop either chronic HBV infection or, rarely, fulminant hepatitis. [1] The role of antivirals, such as lamivudine or entecavir, in acute HBV infection, has not been evaluated in controlled trials.

Lamivudine administration shows an antiviral efficacy in patients with HbeAg positive and HbeAg negative.

Among the patients with acute infection there are elderly subjects with severe course and severe outcome of HBV hepatitis.

A logical hypothesis for these patients is that a rapid decrease in the HBVDNA levels through the use of antiviral agents could result in a less intense host response against HBV virus.

Today, increased knowledge of the virological and immunological events to HBV infection permits to define the

mechanisms involved in viral clearance, persistence and disease severity.

Outcome of infection and the pathogenesis of liver disease are determined by viral and host factors. [1]

The impact of viral load on antiviral T-cell responses has been precisely characterized in animal models of viral infections: a sustained presence of viral antigens leads to virus specific T cell deletion [2].

In HBV infection the frequency and function of circulating and intrahepatic HBV specific CD8 T cells is inversely proportional to the level of HBV-DNA. [3]

The immunological defects could be proportional to the level of HBV infection and inhibition of viral replication and through antiviral treatment it is possible to obtain a partial restoration of HBV specific T cell immunity [4,5] which is inadequate in elderly patients. [6]

Moreover recent studies have provided evidence that a population of specialized T cells are able to regulate the immune response.

These cells reside mainly within a minor population of CD4 cells that express the phenotype marker CD 25. [7]

They have been shown to suppress immunological responses against self and foreign antigens through suppressive cytokines.

It is possible that CD4+ CD25+ T cells are responsible for the weak HBV specific T cell response in HBV infection and may inhibit the expansion and function of HBV specific CD8 T cells precluding HBV clearance.

Some authors showed that the frequency of CD4+ CD25+ T cells positively correlate with HBVDNA load. [6]

This result suggests that an increased level of T regs may be associated with a negative immune response, leading to poor viral clearance.

Our hypothesis is that a decrease of HBV DNA load, determined by lamivudine treatment, determines a downregulation of T regs.

T regs could be able suppressing the population and gamma IFN production, mediated by HbsAg.

In summary, our findings suggest that it could be possible a marked increase in circulating Tregs in elderly patients with HBV infection.

Taken together these data, it is possible that antiviral drugs might be useful in a selected group of old patients in which HBV may cause a severe form of acute hepatitis, caused by a decline of immunity and by the frequent presence of other morbidities.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

GM performed the study and wrote the article.

AP and OP designed the study and reviewed the article.

CS, PB and VR analyzed the clinical data for the study.

References

1. Lok AS, McMahon BJ: **Chronic hepatitis B.** *Hepatology* 2001, **34**:1225-41.
2. Maini HK, Boni C, Oggs GS: **Direct ex vivo analysis of hepatitis B virus specific CD8+ T cells....with the control of infection.** *Gastroenterology* 1999, **117**:1386-96.
3. Dusheinko G: **Hepatitis B.** *J Hepatology* 2003, **44**:948-89.
4. Boni C, Penna A, Bertolotti A, Lamona V, Rapti I, Cavalli A, Cerioni S, Panebianco R, Jenkins J, Ferrari C: **Transient restoration R, of anti-viral T cell responses induced by lamivudine therapy in chronic hepatitis B.** *J Hepatol* 2003, **39**:595-605.
5. Rigopoulou EI, Suri D, Chokshi S, Mullerova I, Rice S, Tedder RS, Williams R, Naoumov NV: **Lamivudine plus interleukin-12 combination therapy in chronic hepatitis B: antiviral and immunological activity.** *Hepatology* 2005, **42**:1028-36.
6. Suvas S, Kumaraguru U, Pack CD, Lee S, Rouse BT: **CD4+CD25+ T cells regulate virus-specific primary and memory CD8+ T cell responses.** *J Exp Med* 2003, **198**:889-901.
7. Guidotti LG, Chisari PV: **Noncytolytic control of viral infections by the innate and adaptive immune response.** *Annu Rev immunol* 2001, **19**:65-91.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

