

**Prevalence of Cervical Neoplastic Lesions and Human Papilloma Virus Infection in Egypt:
National Cervical Cancer Screening Project**

Howayda Abd El All (1), Amany Refaat (2) and Khadiga Dandash (2)

1) Department of Pathology and 2) Department of Community Medicine,
Faculty of Medicine Suez Canal University, Ismailia, Egypt

Corresponding author

Howayda Abd El All: Department of Pathology, Faculty of Medicine, Suez Canal University,
Ismailia, Egypt

Email: howayda@link.net

Abstract

Background: Data from Egyptian studies provide widely varying estimates on the prevalence of pre-malignant and malignant cervical abnormalities and human papilloma virus (HPVs) infection. Objectives: To define the prevalence and risk factors of pre-invasive and invasive cervical cancer (cacx), a community based full-scale cross sectional, household survey including 5453 women aged between 35 and 60 years was conducted. Methodology: Initially, conventional Papanicolaou (Pap) smears were evaluated using the Bethesda system (TBS), followed by colposcopic guided biopsy (CGB) for all epithelial abnormalities (EA). In a third step, HPV was evaluated by in-situ hybridization (ISH) using first the broad spectrum HPV probe recognizing HPVs 6, 11, 16, 18, 30, 31, 35, 45, 51 and 52 followed by subtyping with probes 6/11, 16/18 and 31/33. Lastly, unequivocal cases were immunostained for herpes simplex type-2 (HSV-2), cytomegalovirus (CMV), and human immunodeficiency virus (HIV). Results: EA representing 7.8% (424/5453), were categorized into atypical squamous cell of undetermined significance (ASCUS) (34.4%),

atypical glandular cell of undetermined significance (AGCUS) (15.3 %), combined ASCUS and AGCUS (3.1%), low grade squamous intraepithelial lesions (SIL) (41.0%), high grade SIL (5.2%) and invasive lesions (1%). CGB of EA (n=281) showed non neoplastic lesions (12.8%), atypical squamous metaplasia (ASM) (19.2%), cervical intraepithelial neoplasia (CIN) (51.6%), endocervical lesions (5.2 %), combined squamous and endocervical lesions (10.0%), invasive squamous cell carcinoma (SCC) (0.02%) and extranodal marginal zone B cell lymphoma (MZBCL) (0.02%). The overall predictive value of cytology was 87%. On histological basis, HPVs were present in 94.3% of squamous lesions while it was difficult to be identified in endocervical ones. The prevalence of HPV in the population studied was 2.6%, with HPV 6/11, 16/18 and 31/33 representing 0.4%, 1.4% and 0.7% respectively. Multiple HPVs infections were identified in 0.02%. Conclusion: Pre-invasive and invasive cervical carcinoma represent 2.7% and 0.04% respectively in Egyptian women. HPV mostly 16/18 is a major risk factor ($p < 0.001$), frequently associated with mixed infections ($p < 0.001$) and bilharzial infestation ($p < 0.001$).

Background

In developed countries, with effective and extensive screening with Pap smears, it is usually possible to identify and treat asymptomatic precursor lesions of cacx, making it nearly 100% preventable (1). However, five out of six women with cacx live in developing countries, and 80% of them are diagnosed at advanced stages (2). Data from Egyptian studies provide widely varying estimates on the prevalence of pre-invasive cervical lesions ranging from 1% (3) to 8% (4). Invasive lesions, represented 59.58% of all female genital tract malignancy according to Egyptian National Cancer Institute data (5), and a recent monograph from the Egyptian National Cancer Registry of seven cancer centres of the Ministry of Health and Population, reported a range from 0.12% to 0.77% depending on geographical regions being more prevalent in lower Egypt (6). To be stressed here, that most patients presented at late incurable clinical stages (7, 8). Previous small

hospital based series confirmed the incorporation of HPV genome in invasive lesions (8, 9). In addition schistosomiasis (8,10), Chlamydia trachomatis (Chlamydia T) (11), has been found to be risk factors in Egyptian studies.

In the present work we looked for possible risk factors in the Egyptian population, especially HPV, as a primary risk factor for cacx (12, 13, 14, 15, 16, 17). Other viruses have been implicated as risk factors, such as HSV-2 (16, 18), CMV (16), and HIV (17, 19). Sexual activity includes having multiple sexual partners, sex with a promiscuous partner, history of sexually transmitted infections and sexual intercourse at a young age (20, 21, 22). Bacterial vaginosis (23), Chlamydia T (11, 24), schistosomiasis (8, 10, 25), poor socioeconomic and low educational status (26, 27, 28, 29), risky jobs (30) and poor nutritional status ((31) have been also regarded as risk factors.

This study was the first one to be conducted in Egypt at a National level as community based full-scale cross sectional survey. The main objective was to assess the prevalence of pre-invasive and invasive cacx among Egyptian women aged between 35 and 60 years. Specifically it aimed at (1) assessing the prevalence of biopsy-confirmed squamous, endocervical lesions and invasive cancer, (2) identifying the preventable risk factors and (3) exploring the knowledge and practices of the target population toward cacx screening. Therefore representative data on this problem should permit a better evaluation of possible alternative approaches for the design of a national cacx prevention program.

Materials and Methodols

Sampling

Sampling design and target population

The prevalence rates used for sample size calculations were chosen based on available data from Egyptian studies and results of recent computer modelling. A computer simulation model of incidence of cacx, based on a low risk population in the United States, suggested that the prevalence of SIL, in a previously unscreened population would be 2-3% among women 35 years old and above. The sample size estimates assume simple random sampling equal to 3012 eligible women. Assuming a design effect of 2 due to cluster sampling, the sample size needs to be doubled to be 6000. The target population were ever-married women aged 35 to 60 years. Women who were pregnant or menstruating at time of tests were excluded.

Selection of study population

A multi-stage sampling process was used, based on a sampling frame from the Egypt Demographic Health Survey (DHS) 2000, (32). A sample of different regions of Egypt was made using data from the Human Development Index, which is a composite index of education, health status and living standard. Ten governorates were chosen for the survey, representing urban/rural regions including Lower Egypt (LE), Upper Egypt (UE) and Urban governorates. From each of the sample governorates, 60 clusters of 230 households were selected that were expected to include 125 eligible women. Allowing for refusals would give about 100 women per cluster.

Ethical considerations

The study protocol was approved by the Ministry of Health and Population Ethical Review Board. All women were informed about the procedures of the study, the possibility of abnormalities being found, including cacx and that further invasive procedures may be necessary. Consents were obtained from the women before participation.

Data collection

Questionnaire

A total of 153 questions collected data about women's background and relevant risk factors including 1- socioeconomic and general characteristics of women and their husbands, living conditions, lifestyle, marital status, menstrual and obstetric history; and use of family planning, 2- current general health problems with special emphasis on schistosomiasis, 3- gynecological health problems including personal and family history of cancer or other cancers and women and husbands' history of genital tract infections.

Clinical and gynecological examinations

General clinical examinations and urine laboratory test for schistosomiasis by spotting positive ova were carried out. Problems such as cystocele, rectocele, adnexal tenderness, or masses, uterine prolapse, cervical polyps, warts and cervical discharge, were noted and evaluated. Pap smears were taken.

Pathological study

Cytopathological examination

Samples from the ectocervix and endocervix were collected using Ayer's spatula and endocervical brush and fixed onto labelled glass slides with cytospray. The cells were classified according to TBS (33, 34, 35). Strict criteria were used to qualify a lesion as HPV related (36), ASCUS (36, 37), AGCUS (38), EGD (39) and to discriminate AGCUS from neoplastic endocervical cells (40) (table 1).

Colposcopic-guided biopsy and Histopathology

All EA were biopsied in an attempt to evaluate the prevalence of low / high grade dysplastic changes. The received biopsies were left for 24 hours in neutral buffered formalin. Tissue embedding and processing were followed by hematoxylin and eosin (H&E) staining. The presence of bilharzial infestation was evaluated in tissue sections. Serial sections from each biopsy were classified according to the World Health Organization (WHO) for cervical neoplasia (41). Here

also strict criteria were used for histopathologic evaluation. To qualify a lesion as HPV-related, a scoring system was used for which a minimum of 6 points out of 10 were needed (36). Data not fulfilling these criteria were classified as ASM (table 1).

In-situ hybridization for HPVs

ISH for HPVs was effectuated for epithelial lesions starting ASM. The materials used are summarized in table 2. Initially, HPVs were screened with the wide spectrum HPV probe recognizing genomic DNAs of HPV types 6, 11, 16, 18, 30, 31, 31, 33, 35, 45, 51 and 52. Further subtyping with HPV 6/11, 16/18 and 31/33 were performed for cases positive for pan HPV and cases negative/unsatisfactory for pan HPV but with frank HPV related lesions on conventional H&E biopsies. Positive control tissues infected with HPV were included in every ISH procedure and the technique was evaluated according to the result of this positive control. The steps for ISH are followed according to the manufacturer instructions. In brief, following deparaffinization, tissue digestion with 250-500 μ l of pepsin HCl, were applied to tissue sections and the latter were incubated in 37°C oven for 10 minutes, followed by slides wash in distilled water. Denaturation and hybridization started by applying enough drops of the probe to tissue sections, coverslipping, laying the slide on pre-warmed slide warmer, and incubation in 37°C oven for 60 minutes. Slides were immersed in Tris buffer to remove covers, washed and immersed in warm stringent wash, incubated in 37°C oven for 30 minutes, and then washed in Tris buffer. For revelation, streptavidin alkaline phosphatase was applied to sections for 20 minutes. Slides were washed in Tris buffer twice. BCIP/NBT were applied to sections and incubated in dark for 60 minutes, followed by wash in Tris buffer twice. Slides were lastly washed distilled water and mounted using aqueous media. Positive signals were seen as dark blue dots in the affected nuclei.

Immunohistochemistry

Immunohistochemistry (IHC) was effectuated for equivocal cases negative for HPV and cases with histological suspicion for infections with HSV-2 or CMV. In these instances, HSV-2, early and late CMV antigens were assessed. For other viral lesions not proven cytologically, histologically or by IHC to be HSV-2 or CMV infection, p24 capsid proteins for HIV was used. In addition, cytokeratin, leucocyte common antigen and CD79a were used to diagnose an extranodal MZBCL. The materials used are summarized in table 2. The steps of IHC are followed according to the manufacturer instructions. All incubations were performed at room temperature. Tris buffer was used for washing. In brief, following deparaffinization, the endogenous peroxidase activity was quenched by 0.3% hydrogen peroxide for 5 minutes followed by washing. Primary antibodies diluted at 1/50, were incubated for 30 minutes, then slides were rinsed in successive buffer baths. The revelation was done with LSAB-2 detection kit. Finally, diaminobenzidine tetrachloride (DAB) was applied for 5 minutes. Slides were counterstained in Harris haematoxylin (Hx), dehydrated, cleared in xylene and coverslipped. Tissues infected with HSV-2, CMV and p24 were used as positive control while replacement of the antibodies by Tris buffer was used as negative control for the procedure. IHC results were evaluated either as positive or negative.

Statistical analysis

Lesions were grouped according to histological results into: 1- normal without epithelial changes, 2- ASM, 3- low grade lesions including CINI with or without HPV and EGD alone or combined with either of ASM or CINI, 4- High grade lesions including CINII, CINIII/CIS alone or combined with EGD and adenocarcinoma in situ (AIS), and 5- Invasive lesions. A cut-off point of 45 years was used to categorize two aged groups 35-45 and >45-60 years. SPSS version 9 was used for statistical analysis. Descriptive analysis was made of demographic, cytology and follow-up CGB data. Cross-tabulations with significance tests and Odd Ratio (OR) with 95% confidence interval

(CI) were made. Multivariate analysis was carried out to estimate the main determinant risk factors for cacx in the study population.

Quality control

A pilot study was carried out to test methods of obtaining community cooperation and reactions of respondents to the research procedures and their desire to participate. All staff members were trained to ensure standardization of methods. A written operational manual explaining in details how to perform Pap smear and CGB and all staining steps was prepared. Random re-screening of 20% of the Pap smear and all EA was done for every cluster, in addition to the strict criteria used for evaluating abnormal lesions.

Results

History cervical cancer and awareness of the risk factors

Twelve women (0.2%) had a female relative diagnosed as cacx. Less than 2%, mostly from urban areas, had ever had Pap smear with normal results in 77% of them. Only three women mentioned having dysplastic/atypical changes before, however these findings were not documented by cytopathological reports and their Pap smears were normal in the present study. Only 2% could identify smoking, hormones, and infection as risk factors but not HPV.

Cytopathology

The total number of Pap smears studied was 5658. However, 5453 women with complete data were included in the study (97%). Unsatisfactory results were present in 195 smears (3%), mostly due to absence of endocervical/transformation zone and poor fixation in 66.15% and 16.92% respectively, and were more pronounced in the first three governorates. Two third of the women had inflammatory changes (63%), while epithelial abnormalities were identified among 8%.

Inflammatory and Reparative Changes

All the inflammatory smears were associated with reparative changes including reserve cells hyperplasia, early immature squamous metaplastic cells, cells merging to maturation and fully mature cells. This process was more pronounced in association with IUD devices. Three smears showed mature reparative changes only, probably representing post inflammatory squamous metaplasia. The most common infection was bacterial (49%) (figure 1).

Epithelial abnormalities according to TBS

Epithelial changes were found in 424 women (8%), more commonly among LE women (9%) and less commonly in rural UE (6%); one-third of these cases were among urban governorates residents ($p < 0.005$). The most common type was LGSIL including HPV infection (41%) (figure 2). HGSIL was found in 22 women (5%). Cytological features highly suggestive of invasive carcinoma were diagnosed in four cases. Table 3 illustrates the EA.

Histological evaluation of CGB

CGB was recommended for all women with EA (424 ones). A total of 281 ones (66%) were available. Unsatisfactory biopsies were encountered in 30 cases (10.67%); the most common cause was absence of endocervical tissue in a lesion previously diagnosed as AGCUS (62%), and was more obvious in earlier governorates. The histological results are summarized in table 4. The strict criteria used for evaluating HPV's histologically enabled us to categorize squamous lesions with or without HPV infection (table 5). However, there were no valid criteria for identification of HPV's in endocervical lesions, making ISH a premium for diagnosis of HPV's infected endocervical lesions. The overall predictive value of cervical cytology was 87% with values of 90.10%, 86.68%, 80% and 78.13% for LGSIL, ASCUS, HGSIL and AGCUS respectively. **Figures 3, 4 and 5** illustrate cyto-histologic correlation of some cases.

IHC results

Only one case cytologically diagnosed as ASCUS, was positive for HSV-2. All suspicious cases tested for CMV and HIV infections were negative by IHC. Accidentally, one case positive for leukocyte common antigen and CD79a was classified as extranodal MZBCL.

Prevalence of HPV and cervical lesions

ISH using pan HPV probe tested on 217 cases, was positive among 66%, negative in 29% and non significant in 5% of cases (figures 6, 7 and 8). Interestingly, 12.5% (6/48) of ASM were positive. The prevalence of cervical lesions and HPV infection in the population is summarized in table 6.

Risk factors of cervical lesions

Risk factors of ASM were infection with HPV especially subtypes 16/18 and 31/33 and ever usage of vaginal contraceptives ($p < 0.0001$ for each). The risk factors for low grade CIN are still menstruating ($p < 0.001$), working ($p < 0.0001$), HPV infection with the three probes studied 6/11, 16/18 and 31/33 ($p < 0.000$ for each), while for high grade CIN infection with HPV mainly subtypes 16/18 and 31/33 ($p < 0.001$) was the common threat. Multiple regression analysis revealed that being a working woman (OR 2.0, 95% CI 1.3-3.1) and still menstruating (OR 1.5, 95% CI 1.1-2.1) were the main risk factors.

HPVs infection were more pronounced in younger age, actually married, still menstruating and ever used hormonal or vaginal contraceptives ($p < 0.05$ for each), and working especially in risky jobs ($p < 0.0001$). Applying multiple regression analysis revealed that being a working woman (OR 2.0, 95% CI 1.3-3.2), still menstruating (OR 1.6, 95% CI 1.1-2.3) and women who ever had hormonal contraception (OR 1.5, 95% CI 1.1-2.1) were the main determinants for HPV infection. HPV infection seemed to be more among women whose husbands ever had another wife or travelled before, however this finding was not statistically significant.

Discussion

Awareness of HPV as major risk factor for cacx was very low among the studied population while smoking, hormones, and infections were identified as main risk factors. Previous studies approaching this point came out with the same conclusion of ignorance of association between HPV and cacx (42, 43). As expected, only 1.5% of studied women, mostly from urban areas, had Pap smear. Three of them mentioned, without documents, having results consistent with invasive lesions. However, their cytological results were normal in this study. This important point highlights the absence of health culture in the population. Studies from other countries, reported that only 5% of women in low and middle income countries received screening, usually in private clinics and few urban settings, contrary to high-income countries where up to 70% of women were screened (1, 44).

ASM cases were positive for HPV 16/18 and 31/33. ASM, a poorly characterized cervical lesion with uncertain biological and clinical significance, shares some but not all morphological features of SIL (45, 46). It has been shown that HPV positive ASM biopsies were significantly more likely to have concurrent or subsequent diagnosis of HSIL than HPV negative ones (45, 47).

Unfortunately, the actual study is limited by the fact that it covers a single period and had no opportunity to follow up ASM HPV positive cases. The rationale for inclusion of ASM is based on the fact that recent studies found HPV's genome incorporated in normal squamous epithelium (22, 48, 49). However, due to financial limitations, normal tissues were not included, to evaluate the presence/absence of HPV's.

This community based study confirm previous hospital based one reporting low prevalence rate for CIN; analysis of 4458 patients showed a prevalence of 0.36%, 0.23 % and 0.12 % for CINI, CINII and CINII respectively (50). In addition, the Middle East Cancer Consortium reported 0.027% prevalence rate for Egyptian women (51). The prevalence of EA and HPV infections in Egypt is comparable to other geographical areas in Muslim and Middle East countries. In Saudi Arabia, EA

represented 3.14% out of 3088 screened women with ASCUS, LGSIL, HGSIL, invasive SCC, AGCUS and adenocarcinoma represented 0.45%, 0.93%, 0.55%, 0.13%, 0.13% and 0.03% respectively (52). In Jordan a prevalence of 0.026% has been reported (51). In Lebanon, a study conducted on 1,026 women revealed a prevalence of 4.9% for HPV with type 16 representing 3% (53). In Morocco, 70.5% of invasive carcinoma cases were HPV positive, 34.88% cases had HPV16, and 15.5% cases had HPV18 (54). In Iran, controversial data were reported; 73.9% of HPV positive cacx contained HPV16 (55), while in another study HPV16 was identified in 26.7% and none of the samples were positive for HPV18 (56).

Studies showed significant geographic variation in the prevalence of oncogenic viral types in cervical lesions with roughly half of all cacx worldwide containing HPV 16. Other important high risk types are HPV 18, 45, and 31. Less prevalent high risk types include HPV 26, 33, 35, 39, 51, 52, 56, 58, 59, 68, 73, and W13b (49). In Africa, the most common subtype is HPV 16 followed by 45, 18, 31 and 33 (57), a point concordant with the actual work. Unskilled workers were at greater risk for HPV infection and other EA. These women were the highest to have husbands with history of genital infection, showed vaginal discharge by examination, reported history of infection and higher sexual activity. HPV infection seemed to be more among women whose husbands ever had another wife or travelled before. In addition, one fourth of unskilled workers had husbands who ever travelled before or married another wife. In spite that the majority of women had married once, 15% of unskilled workers had married more than once. It is clear that the marital/sexual practices of unskilled workers were the determinants of their higher susceptibility to HPV and CIN. The socio-economic characteristics of unskilled workers confirm previous studies where associations have been found between low socioeconomic status (1, 28, 26,), multiple sexual partners (58, 59, 60, 61) and cacx. Another risk factor, smoking, is added to unskilled workers, as 83% of them were exposed to smoking at home and work. Higher prevalence for high-risk HPV types was significantly associated with cigarette smokers (62). Steroid contraception has been

postulated to be one mechanism whereby HPV exerts its oncogenic effect on cervical tissue especially among long-term users (63), a point concordant with the present study, mostly noted in one half of unskilled workers.

Our data on schistosomiasis as risk factor for the development of cacx confirm previous studies in Egypt (8, 10) and elsewhere (25). However, *T. vaginalis* previously implicated to have direct relation to cacx in the Egyptian population (11) were not established in the actual work. Moreover, early marriage or early sexual relations, was not significantly associated with HPV as Egyptian women start sexual relations with marriage. Other studies confirmed that women who have their first intercourse at an early age are at high risk for HPV infection and cacx (58, 59, 64), a point not tested as Egyptian women start sexual relations with marriage. Being uncircumcised male was found to be a high risk factor (61), which was not evaluated due to universality of male circumcision among Egyptian men both Moslem and Christians.

The study is limited by the fact that it covers a single period and had no opportunity to examine trends over time. Three obstacles reduced the number of satisfactory smears, noted mostly at the beginning of the study, implying the importance of training. These were, smears not interesting the squamo-columnar junction, inadequate fixation and hemorrhagic smears resulting from traumatization of the mucosa by the cytobrush. Inadequate biopsies were encountered whenever they were performed in health units using portable colposcope. In addition, women of higher socioeconomic group refusal to participate may have led to missing the chance to describe the pattern of cervical lesions among this group. The drop out of women for CGB was due to their refusal to participate in spite of counselling. Concerning HPV's ISH, two points should be discussed. The first was fixation more than 24 hours in UE specimens, a point that may explain the negativity for all HPV's probes even with the presence of frank HPV's in biopsies. The second was

the limited number of tested probes compared to the known HPV's subtypes. Cultural sensitivity for sexual relationships in the Egyptian community in general and among women specifically, made it difficult to get detailed information even with using probing questions for assessing sexually transmitted infections.

In conclusion, this study is the first one at a national level describing the characteristics of women with cervical epithelial abnormalities. They were mostly of middle income, married, with three and more children, mostly uneducated and not working. The awareness of the Egyptian women by risk factors or symptoms of different reproductive health problems are extremely low. Prevalence of CIN and invasive lesions was 3.1% and 0.04%, while the prevalence of HPV's was 2.6% and was positive in 94.3% of cervical lesions confirming that it is the main causing agent. The study recommends introducing cax screening for all women at least once every 10 years for women with normal cytological findings, and yearly for three successive years for inflammatory changes. CGB should be performed to all EA to avoid losing patients and subsequently tumor progression. In addition, raising awareness of Egyptian women on reproductive health and risk factors of cax through specially designed health communication programs is mandatory.

Contributions

HA was the Principal Investigator for Pathology, AR was the Principal Investigator for Data Management and KD was the Principal Investigator for Field Work.

Acknowledgement and Collaborators

The study was carried out by the Faculty of Medicine Suez Canal University in collaboration with the Research Management Unit of the National Population Council and sponsored by the United States Agency for International Development (USAID/Cairo). The authors are grateful to other members of the Faculty participating in the Project; the Gynecology Team, Professor Galal Lotfy

and Rasha el Saied, Marium Lotfy, Abeer Bahaa, Amal Saied, Nahed Amin, Mona Fathy, Thuria Ahmed, Mona Galal Eldine, Soheir Abdelghafar and Kawser abdelazeim, Assistant Lecturers of Obstetrics and Gynecology; the colposcopy team, Mohamed Shaban, Yaser Habash, Mosaad Abdelhamid, Mohamed Abouelsood , Mohey Mohamed; the pathology team, Maha Rashwan and Hala Abd El Dayem Assistant Lecturers and Magdy El Korachy, Demonstrator; field work team, Aymen Farghaly, Rasha Farouq, Hatem Salam and Yasser Youssef, Community Medicine Demonstrators and Tamer Abdel Fatah, Microbiology. The cooperative assistance of the cytopathologists Professors Leila Nabegh, Magda Abd El Salam and Zeinab Chegab Eldine and the Associate Professor Rageya Ahmed at Ain Shams University is appreciated. We are grateful to the enormous great effort of the Pathology Technicians Fatouh Arafa, Mohammed El Seify, Sayed Chaheen, Mohammed Saeed, Nahla Abd El Reheem, Cherine Farouok, Mohammed El Cherbeeney, Mr Sayed Gareeb and Zeinab Ahmed. The secretarial assistance of Rabaab Metwally was very helpfull. Special thanks to the health units' directors, interviewers, listing personal, social workers and nurses. This study wouldn't take place without the help and encouragement of our ex Dean Professor Mohammed Abd El Kafy and our actual Dean Professor Mohamed Shoeir.

Abbreviations

AGCUS: atypical glandular cells of undetermined significance, AIS: adenocarcinoma in situ, ASCUS: atypical squamous cells of undetermined significance, ASM: atypical squamous metaplasia, Cax: cancer cervix, Chlamydia T: Chlamydia trachomatis, CIN: cervical intraepithelial neoplasia, EA: epithelial abnormalities, EGD: endocervical glandular dysplasia, HGSIL: high grade squamous intraepithelial lesion, HPVs: human papilloma viruses, IHC: Immunohistochemistry. ISH: in situ hybridization, LGSIL: low grade squamous intraepithelial lesion, Pap smears: Papanicolaou smears, TBS: the Bethesda System, SCC: squamous cell carcinoma.

References

1. Murray CJL, Lopez AD. Global health statistics. Boston, Harvard School of Public Health on behalf of WHO and the World Bank, 1996.
2. Stewart BW, Kleihues P. Cancers of the female reproductive tract. In, WHO Cancer Report. Lyon: IARC pp 215- 222, 2003.
3. El Mosselhy MH, Moselhy MM, Amin A, Ouda-Pasha E, Abdel Hamid, Hussein MH. Prevalence of reproductive tract infections among married women of reproductive age. National Population Council/El Galaa Teaching Hospital, 1998.
4. Younis N, Khattab H, Zurayk H, el-Mouelhy M, Amin MF, Farag AM. A community study of gynecological and related morbidities in rural Egypt. *Stud Fam Plann* 24:175-186, 1993.
5. Mokhtar N. Cancer Pathology Registry 1985-1989. Department of Pathology. NCI, 1991.
6. National Cancer Registry. The National Cancer Registry Newsletter. Ministry of Health and Population. 2002.
7. Abd El All H, Nada G, Awaad HK, El Ayaat A. Natural killer cells in human cervical carcinoma. Master Degree Thesis, Suez Canal University, Ismailia, Egypt, 1987.
8. Abd El All H, Prade M, Khatab T, Awaad HK, Ghaly AF. Immunological study of NK cells in cervical carcinoma. Ph.D. Thesis, Faculty of Medicine, Suez Canal University, Ismailia, Egypt, 1992.
9. Ahmed MI, Salahy EE, Fayed ST, El-Hefnawy NG, Khalifa A. Human papillomavirus infection among Egyptian females with cervical carcinoma: relationship to spontaneous apoptosis and TNF-alpha. *Clin Biochem.* 34:491-498, 2001.
10. Youssef AF, Fayad MM, Shafeek AED. Bilharziasis of the cervix uteri. *BJOG* 77: 847-851, 1970.

11. Sayed el-Ahl SA, el-Wakil HS, Kamel NM, Mahmoud MS. A preliminary study on the relationship between *Trichomonas vaginalis* and cervical cancer in Egyptian women. *J Egypt Soc Parasitol* 32: 167-178, 2002.
12. Zur Hausen H. Intracellular surveillance of persisting viral infections. *Lancet* 30: 489-491, 1986.
13. Zur Hausen H. Papilloma viruses in anogenital cancer as a model to understand the role of viruses in human cancers. *Cancer Res* 49: 4677-851, 1989.
14. Zur Hausen H. Papillomavirus infections-a major cause of human cancers. *Biochim Biophys Acta*. 1288: 55-78, 1996.
15. Zur Hausen H. Papillomaviruses causing cancer: evasion from host-cell control in early events in carcinogenesis. *J Natl Cancer Inst* 92: 690-698, 2000.
16. Han CP, Tsao YP, Sun CA, Ng HT, Chen SL. Human papillomavirus, cytomegalovirus and herpes simplex virus infections for cervical cancer in Taiwan. *Cancer Lett* 120: 217-221, 1997.
17. Rezza G, Giuliani M, Branca M, Benedetto A, Migliore G, Garbuglia, D'Ubaldo C, Pezzotti P, Capiello G, Pomponi-Formiconi D, Suligoi B, Schiesari G, Giacomini G. Determinants of squamous intraepithelial lesions (SIL) on Pap smear: the role of HPV infection and of HIV-1-induced immunosuppression. DIANAIDS Collaborative Study Group. *Eur J Epidemiol* 13: 937- 943, 1997.
18. Viikki M, Pukkala E, Nieminen P, Hakama M. Gynaecological infections as risk determinants of subsequent cervical neoplasia. *Acta Oncol* 39: 71-75, 2000.
19. Palefsky JM, Minkoff H, Kalish LA, Levine A, Sacks HS, Garcia P, Young M, Melnick S, Miotti P, Birk P. Cervicovaginal human papillomavirus infection in human immunodeficiency virus-1 (HIV)-positive and high-risk HIV-negative women. *J Natl Cancer Inst* 91: 226-236, 1999.

20. Lovejoy NC. Precancerous and cancerous cervical lesions: the multicultural "male" risk factor. *Oncol Nurs Forum* 21: 497- 504, 1992.
21. Bosch FX, Castellsague X, Munoz N, de Sanjose S, Ghaffari AM, Gonzalez LC, Gili M, Izarzugaza I, Viladiu P, Navarro C, Vergara A, Ascunce N, Guerrero E, Shah KV. Male sexual behavior and human papillomavirus DNA: key risk factors for cervical cancer in Spain. *J Natl Cancer Inst* 88: 1060-1067, 1996.
22. Bosch FX, Lorincz A, Munoz N, Meijer CJ, Shah KV. The causal relation between human papillomavirus and cervical cancer. *J Clin Pathol.* 55: 244-265, 2002.
23. Castle PE, Wacholder S, Lorincz AT, Scott DR, Sherman ME, Glass AG, Rush BB, Schussler JE, Schiffman M. A prospective study of high-grade cervical neoplasia risk among human papillomavirus-infected women. *J Natl Cancer Inst* 94: 1406-1414, 2002.
24. Markowska J, Fischer N, Fischer Z, Warchol JB. Chlamydia trachomatis infection in women with CIN and invasive uterine cervix cancer. Significance of hormonal status. *Eur J Gynaecol Oncol.*23: 511- 513, 2002.
25. Moubayed P, Ziehe A, Peters J, Mwakyoma H, Schmidt D. Carcinoma of the uterine cervix associated with schistosomiasis and induced by human papillomaviruses. *Int J Gynaecol Obstet* 49:175-179, 1995.
26. de Sanjose S, Bosch FX, Munoz N, Tafur L, Gili M, Izarzugaza I, Izquierdo A, Navarro C, Vergara A, Munoz MT, Ascunce N, Shah KV. Socioeconomic differences in cervical cancer: two case-control studies in Colombia and Spain. *Am J Public Health* 86: 1532-1538, 1996.
27. de Sanjosé S, Bosch FX, Muñoz N, Shah K. Social differences in sexual behaviour and cervical cancer. *IARC Sci Publ* 138: 309-317, 1997.
28. Mackillop WJ, Zhang-Salomons J, Boyd CJ, Groome PA. Associations between community income and cancer incidence in Canada and the United States. *Cancer* 89: 901-912, 2000.

29. Wellensiek N, Moodley M, Moodley J, Nkwanyana N. Knowledge of cervical cancer screening and use of cervical screening facilities among women from various socioeconomic backgrounds in Durban, Kwazulu Natal, South Africa. *Int J Gynecol Cancer* 12: 376- 382, 2002.
30. Weiderpass E, Pukkala E, Vasama-Neuvonen K, Kauppinen T, Vainio H, Paakkulainen H, Boffetta P, Partanen T. Occupational exposures and cancers of the endometrium and cervix uteri in Finland. *Am J Ind Med.* 39: 572-580, 2001.
31. Ramaswamy G, Krishnamoorthy L. Serum carotene, vitamin A, and vitamin C levels in breast cancer and cancer of the uterine cervix. *Nutr Cancer* 25: 173-177, 1996.
32. El Zanaty F, Way A. EDHS 2000: Egypt Demographic and Health Survey, Ministry of Health and Population. National Population Council. 2001.
33. Solomon D. The Bethesda system for reporting cervical/vaginal cytologic diagnoses. *Acta Cytol* 33: 567- 574, 1989.
34. Solomon D. The Bethesda System for reporting cervical/vaginal cytologic diagnosis: an overview. *Int J Gynecol Pathol.*10: 323- 325, 1991.
35. Solomon D, Davey D, Kurman R, Moriarty A, O'Connor D, Prey M, Raab R, Sherman M, Wilbur D, Wright T, Young N. The 2001 Bethesda System. Terminology for Reporting Results of Cervical Cytology. *JAMA.* 287: 2114- 2119, 2002.
36. Concensus morphologique sur les néoplasies du coluterin. Paris-Tolbiac. *Archives d' Anatomie et de Cytologie Pathologiques* 39:186- 222, 1991.
37. Malik SN, Wilkinson EJ, Drew PA, Bennet BB and Hardt NS. Do qualifiers of ASCUS distinguish between low- and high risk patients? *Acta Cyto* 43: 376- 380, 1999.
38. Morin C, Bairati I, Bouchard C, Fortier M, Roy M, Moore L, Meisels A. Cytologic predictors of cervical intraepithelial neoplasia in women with an ASCUS Pap smear. *Acta Cytol* 44: 576- 586, 2000.

39. Casper GR, Ostor AG, Quinn MA. A clinicopathologic study of glandular dysplasia of the cervix. *Gynecol Oncol.* 64:166-170, 1997.
40. Cenci M, Mancini R, Nofroni I, Vecchione A. Endocervical atypical glandular cells of undetermined significance. I. Morphometric and cytologic characterization of cases that "cannot rule out adenocarcinoma in situ". *Acta Cytol* 44:319-326, 2000.
41. Tavassoli FA, Devilee P. Pathology and Genetics. Tumors of the breast and female genital organs. WHO classification of tumours. Chapter 6, 2003.
42. Pitts M, Clarke T. Human papillomavirus infections and risks of cervical cancer: what do women know? *Health Educ Res* 17: 706- 714, 2002.
43. Waller J, McCaffery K, Forrest S, Szarewski A, Cadman L, Wardle J. Awareness of human papillomavirus among women attending a well woman clinic. *Sex Transm Infect* Aug 79: 320-322, 2003.
44. Miller AB, Nareer S, Fonn S, Brandup-Lukanon A, Rakshanda R, Cronje H, Sankaranarayanan R, Korolcltchouk, Syrjanen K, Singer A, Onsrud M. Report on consensus conference on cervical cancer screening and management. *Int J Gyn Obst*, 86: 440- 447, 2000.
45. Geng L, Connolly DC, Isacson C, Ronnett BM, Cho KR. Atypical immature metaplasia (AIM) of the cervix: is it related to high-grade squamous intraepithelial lesion (HSIL)? *Hum Pathol* 30: 345-351, 1999.
46. Park JJ, Genest DR, Sun D, Crum CP. Atypical immature metaplastic-like proliferations of the cervix: diagnostic reproducibility and viral (HPV) correlates. *Hum Pathol* 30:1161-1165, 1999.
47. Quddus MR, Zhang S, Sung CJ, Liu F, Neves T, Struminsky J, Singer DB. Utility of HPV DNA detection in thin-layer, liquid-based tests with atypical squamous metaplasia. *Acta Cytol* 46: 808- 812, 2002.
48. Manos MM, Kinney WK, Hurley LB, Sherman ME, Shieh-Ngai J, Kurman RJ, Ransley JE, Fetterman BJ, Hartinger JS, McIntosh KM, Pawlick GF, Hiatt RA. Identifying women with

cervical neoplasia: using human papillomavirus DNA testing for equivocal Papanicolaou results. *JAMA* 281:1605-1610, 1999.

49. Muñoz N, Bosch FX, de Sanjosé S, Herrero R, Castellsagué X, Shah KS, Snijders PJF, Chris J.L.M. Meijer CJLM, for the International Agency for Research on Cancer Multicenter Cervical Cancer Study Group. Epidemiologic Classification of Human Papillomavirus Types Associated with Cervical Cancer. *New England J Medicine* 348: 518- 527, 2003.
50. Hammad MM, Jones HW 3rd, Zayed M. Low prevalence of cervical intraepithelial neoplasia among Egyptian females. *Gynecol Oncol.*28: 300- 304, 1987.
51. Komoditi C. Cervical and Corpus Uterine Cancer. P 83-90. In *Cancer Incidence in Four Member Countries (Cyprus, Egypt, Israel, and Jordan) of the Middle East Cancer Consortium (MECC) Compared with US SEER*. Ed: Freedman LS, Edwards BK, Ries LA, Young JL. National Cancer Institute. NIH Pub. No. 06-5873. Bethesda, MD.
52. Altaf FJ. Pattern of cervical smear cytology in the western region of Saudi Arabia. *Annals Saudi Med* 21: 94- 96, 2001.
53. Mroueh AM, Seoud MA, Kaspar HG, Zalloua PA. Prevalence of genital human papillomavirus among Lebanese women. *Eur J Gynaecol Oncol* 23: 429- 432, 2002.
54. Lalaoui K, El Mzibri M, Amrani M, Belabbas MA, Lazo PA. Human papillomavirus DNA in cervical lesions from Morocco and its implications for cancer control. *Clin Microbiol Infect* 9: 144- 148, 2003.
55. Mortazavi S, Zali M, Raoufi M, Nadji M, Kowsarian P, Nowroozi A. The Prevalence of Human Papillomavirus in Cervical Cancer in Iran. *Asian Pac J Cancer Prev.* 3: 69-72, 2003.
56. Farjadian S, Asadi E, Doroudchi M, Dehaghani AS, Tabei SZ, Kumar VP and Ghaderi A. High risk HPV types in southern Iranian patients with cervical cancer. *Pathol Oncol Res.* 9:121-125, 2003.

57. Clifford GM, Smith JS, Plummer M, Muñoz N, Franceschi S. Human Papilloma virus types in invasive carcinoma worldwide: a meta-analysis. *Br J Cancer* 88: 63- 73, 2003.
58. Williams MA, Kenya PR, Mati JK, Thomas DB. Risk factors for invasive cervical cancer in Kenyan women. *Int J Epidemiol* 23: 906-912, 1994.
59. Biswas LN, Manna B, Maiti PK, Sengupta S. Sexual risk factors for cervical cancer among rural Indian women: a case-control study. *Int J Epidemiol* 26: 491- 495, 1997.
60. Murthy NS, Mathews A. Risks for pre-invasive lesions of the cervix. *Eur J Cancer Prev* 9: 5-14, 2000.
61. Svare EI, Kjaer SK, Worm AM, Osterlind A, Meijer CJ, van den Brule AJ: Risk factors for genital HPV DNA in men resemble those found in women: a study of male attendees at a Danish STD clinic. *Sex Transm Infect* 78: 215- 218, 2002.
62. Roteli-Martins CM, Panetta K, Alves VA, Siqueira SA, Syrjanen KJ, Derchain SF. Cigarette smoking and high-risk HPV DNA as predisposing factors for high-grade cervical intraepithelial neoplasia (CIN) in young Brazilian women. *Acta Obstet Gynecol Scand* 77: 678- 682, 1998.
63. Moodley M, Moodley J, Chetty R, Herrington CS. The role of steroid contraceptive hormones in the pathogenesis of invasive cervical cancer: a review. *Int J Gynecol Cancer* 13:103-110, 2003.
64. Kahn JA, Rosenthal SL, Succop PA, Ho GY, Burk RD. The interval between menarche and age of first sexual intercourse as a risk factor for subsequent HPV infection in adolescent and young adult women. *J Pediatr* 141: 718- 723, 2002.

Figures legend

Figure 1: Inflammatory changes.

Figure 2: LGSIL HPV associated showing koilocytic atypia (A), binucleation (B) and dyskeratosis (C) Papanicolaou stain, x 40.

Figure 3: Combined dysplasia, A- ASCUS, the nucleus is twice the size of the left intermediate cell on the right, Pap stain, x40. B- Another area showing nuclear pseudostratification and loss of the normal honey combing, Pap stain, x40. C- A third area showing ASCUS and AGCUS in the same field, Pap stain, x40. D- Glandular colonization by the neoplastic cells and dysplastic endocervical lining, H&E, x10. E- Higher power magnification of the colonized glands showing koilocytes, increased nucleo-cytoplasmic ratio and large nucleolated nuclei, H&E, x40. F- Higher power magnification of the endocervical dysplastic cells, H&E, x40. G- Another field showing colonization of the glands by the koilocytes and residual normal endocervical cells, H&E staining, x40. H- Surface epithelium showing koilocytic atypia, acanthosis, papillomatosis and basal cell hyperplasia, H&E, x40.

Figure 4: A- AGCUS favoring a neoplastic process, Pap stain, x 400. B- AIS, H&E staining, x20.

Figure 5: LCNK SCC, A- Fibres cells (arrow) and koilocytes (arrow head), Pap stain, x 400. B- Marked nuclear pleomorphism and hyperchromasia and heavy neutrophilic infiltrate, Pap stain, x 400. C- Corresponding biopsy: islands of neoplastic cells surrounded by heavy neutrophilic infiltrate, H&E, x10

Figure 6: CINIII, A- H&E, x 20. B- HPV nucleic acid, ISH, pan HPV probe, x 20.

Figure 7: EGD, A- EGD, ISH, pan HPV probe, x10. B- Higher power magnification showing variable degree of intensity with severe intensity in the middle zone, x40.

Figure 8: LCNK SCC A-, H&E, x20. B- HPV nucleic acid in almost all the cells. ISH, pan HPV probe, x 20

Table 1: Cytological and histological criteria used for diagnosis of cervical lesions

Lesions	Criteria for diagnosis	Ref.
Cytologic HPV	<p>Koilocytic atypia</p> <p>Multinucleation (frequently binucleation)</p> <p>Dyskeratosis</p> <p>Parakeratosis</p>	36
ASCUS	<p>Nuclear enlargement approximately twice the size of an immature squamous metaplastic cell</p> <p>Slight increase in nucleo-cytoplasmic ratio</p> <p>Nuclear chromatin clumping</p> <p>Variation in nuclear size and shape including mild hyperchromasia, binucleation, and mild irregularity in the nuclear membranes</p>	37, 38
AGCUS	<p>Nuclear enlargement approximately twice the size of an endocervical cell.</p> <p>Nuclear pseudostratification</p> <p>Mild hyperchromasia</p>	39
Discriminating AGUS from neoplastic endocervical cells	<p>Presence of normal ECCs, singly or in sheets</p> <p>Absence of necrosis</p> <p>Absence of papillary groups</p> <p>Mild anisonucleosis</p> <p>Marked chromatin distribution</p>	39
EGD	<p>Nuclei not cytologically malignant</p> <p>Nuclear hyperchromasia and pseudostratification</p>	40

	Low mitotic figures Absence of papillary formation and cribriform pattern	
Histologic HPV	Koilocytic atypia: 4 points Binucleation: 2 points Dyskeratosis: one point Basal cell hyperplasia: one point Papillomatosis: one point Acanthosis: one point	36

Table 2: Reagents used in the study

Reagent	Source	code
<i>ISH detection kits and probes</i>		
ISH detection system	DakoCytomation	K0601
Wide spectrum HPV	DakoCytomation	Y1404
6/11 HPV	DakoCytomation	Y1411
16/18 HPV	DakoCytomation	Y1412
31/33 HPV	DakoCytomation	Y1413
<i>IHC detection kits and antibodies</i>		
LSAB-2 Detection kit	DakoCytomation	K0680
Herpes simplex virus type 2	DakoCytomation	B0116
CMV early antigens	Novocastra	NCL- CMV- EA
CMV late antigens	Novocastra	NCL- CMV- EA
P 24 capsid protein	Novocastra	NCL-p24

Table 3: Percentage distribution of women by cytological epithelial abnormalities according to place of residence

Epithelial abnormalities	Urban governorates	LE		UE		P	Total	
		Urban	Rural	Urban	Rural		Freq.	%
<i>Atypical</i>						<0.05		
ASCUS	37.2	27.3	35.2	46.3	26.7		146	34.4
AGCUS	14.9	14.5	16.2	29.3	8.0		65	15.3
Combined	1.4	1.8	4.8	2.4	5.3		13	3.1
<i>Squamous lesions</i>	41.2	43.6	37.1	12.2	60.0		174	41.0
LGSIL	4.7	10.9	5.7	7.3	0.0		22	5.2
HGSIL								
<i>Invasive</i>	0.0	1.8	0.0	2.4	0.0		2	0.5
Squamous	0.7	0.0	1.0	0.0	0.0		2	0.5
Endocervical								
Total	148	55	105	41	75	424		

Table 4: Histological results and pan HPV ISH of CGB

Histologic results	Histology		Pan HPV ISH		
	Freq .	%	Negative	Positive	NS
<i>Normal</i>	32	12.8	ND	ND	ND
<i>ASM</i>	48	19.2	39	6	3
<i>CIN</i>					
CINI	111	44.4	16	89	6

CINII	11	4.4	2	9	0
CINIII/CIS	7	2.8	0	7	0
<i>Glandular lesions</i>					
EGD	12	4.8	4	8	0
AIS	2	0.8	0	1	1
<i>Combined dysplasia</i>					
ASM + EGD	4	1.6	2	2	0
CINI+EGD	18	7.2	0	17	1
CINII+EGD	2	0.8	0	2	0
CINIII+EGD	1	0.4	0	1	0
<i>Invasive</i>					
Invasive carcinoma	1	0.4	0	1	0
Lymphoma	1	0.4	ND	ND	ND
Total	250	100	63 (29.0%)	143 (65.9%)	11 (5.1%)
			217 (86.8%)		

ND: not done NS: non significant

Table 5: Histological HPV's infection in neoplastic squamous lesions

	Freq.	Histologic evidence of HPV	%
CIN I	111	104	93.7
CINII	13	12	92.3
CINIII	7	6	85.7
Combined dysplasia	25	25	100.0

Invasive carcinoma	1	1	100.0
Total	157	148	94.3

Table 6: Prevalence of cervical lesions and HPV infection in the Egyptian population

<i>Cervical lesions and HPV infection</i>	Freq.	%
<i>Cervical preinvasive and invasive lesions</i>		
CINI	111	2.07
CIN I+EGG	18	0.07
CINII	13	0.24
CIN II+EGG	2	0.03
CINIII	7	0.13
CINIII + EGD	1	0.02
EGD	12	0.22
AIS	1	0.03
Combined squamous and glandular	25	0.47
SCC	1	0.02
<i>HPV infections</i>		
Pan HPV	143	2.6
HPV 6/11	24	0.4
HPV 16/18	68	1.4
HPV 31/33	37	0.7
	5453	100







Figure 1: Inflammatory changes




A




B




C




A




B




C




D




E




F




G




H




A




B




A




B




C




A




B




A



B



A



B